

CANINE VACCINATION PROTOCOL – 2005

MINIMAL VACCINE USE

**W. Jean Dodds, DVM
HEMOPET
938 Stanford Street
Santa Monica, CA 90403
310-828-4804; Fax 310-828-8251
e-mail: hemopet@hotmail.com**

Note: The following vaccine protocol is offered for those dogs where minimal vaccinations are advisable or desirable. The schedule is one I recommend and should not be interpreted to mean that other protocols recommended by a veterinarian would be less satisfactory. It's a matter of professional judgment and choice.

Age of Pups	Vaccine Type
9 - 10 weeks	Distemper + Parvovirus, MLV (e.g. Intervet Progard Puppy DPV)
14 weeks	Same as above
16 -18 weeks (optional)	Same as above
20 weeks or older, if allowable by law	Rabies
1 year	Distemper + Parvovirus, MLV
1 year	Rabies, killed 3-year product (give 3-4 weeks <i>apart</i> from distemper/parvovirus booster)

Perform vaccine antibody titers for distemper and parvovirus annually thereafter. Vaccinate for rabies virus according to the law, except where circumstances indicate that a written waiver needs to be obtained from the primary care veterinarian. In that case, a rabies antibody titer can also be performed to accompany the waiver request.

W. Jean Dodds, DVM